

1/15




FIG. 1

2/15




FIG. 2

3/15




FIG. 3

4/15




FIG. 4

5/15




FIG. 5

6/15




FIG. 6

7/15

Plasma Concentration of ISIS 3082, 5' C18 Amine (ISIS 9047)
 and 5' Cholesterol (ISIS 8005) Analogs
 in BALB/c Mice After Intravenous Injection




FIG. 7

8/15

ISIS 3082 Tissue Distribution in BALB/c Mice as Percent of Dose at 1,3,8 and 24 Hours After Intravenous Injection




FIG. 8

9/15

ISIS 9047 (5' C18 amine phosphorothioate analog of ISIS 3082)
 Tissue Distribution in BALB/c Mice as Percent of Dose
 at 1, 3, 8 and 24 Hours After Intravenous Injection




FIG. 9

10/15

ISIS 8005 (5' cholesterol analog of ISIS 3082)
Tissue Distribution in BALB/c Mice as Percent of Dose
at 1,3,8 and 24 Hours After Intravenous Injection




FIG. 10

11/15

Inhibition of Liver ICAM-1 mRNA in BALB/c Mice.




FIG. 11

12/15

The 5' - cholesterol 3082 analog - 8005 - has
no effect on c-raf or PECAM-1 mRNA Levels.




FIG. 12

13/15





FIG. 13


14/15

ICAM-1 IHC in LPS Liver



Basal

LPS / No Oligo



LPS / ICAM-1 Cholesterol 5' (8005)

LPS / Scrambled Cholesterol 5' (13293)

FIG. 14

BEST AVAILABLE COPY

15/15

Inhibition of BALB/c Mouse Liver ICAM-1 mRNA with 3' Cholesterol ASO.




FIG. 15